

Estimating Biomass in Coastal *Baccharis pilularis* Dominated Plant Communities

Will Russell¹ and Ryan Tompkins²

¹*Environmental Studies Department
San Jose State University
One Washington Square
San Jose, CA 95192
russell_sjsu@yahoo.com*

²*USDA Forest Service
Plumas National Forest
159 Lawrence Street
Quincy, CA 95971*

ABSTRACT

Communities dominated by *Baccharis pilularis* (coyote brush) are expanding in coastal California, altering fuel load on a landscape scale, yet there is no standard method for estimating biomass in this vegetation type. In an attempt to develop a non-destructive field method for estimating biomass in *Baccharis*-dominated communities, we compared three indirect measures including crown canopy height, basal stem diameter, and leaf area index (LAI); estimated using hemispherical photography. Data were collected on 90 one square meter randomly selected plots in Point Reyes National Seashore and Golden Gate National Recreation Area. Linear regression analysis was used to determine the effectiveness of each predictor. The best single predictor of biomass was crown canopy height with an adjusted R^2 of 0.46 for both sites combined. The linear regression equation developed for biomass versus height predicted 3930 grams per meter in height \pm 430 grams for each square meter on these sites found. Basal stem diameter was determined to be a weak predictor in this vegetation type ($R^2 = 0.17$). Estimated LAI had no predictive value. Multiple linear regressions with biomass versus height and basal area was also tested resulting in a slightly stronger model with an adjusted R^2 of 0.48. Because of its predictive value and ease of implementation, we suggest the use of crown canopy height as the most efficient biomass predictor in this vegetation type.

Keywords: *Baccharis pilularis*, *Coyote brush*, *biomass*, *fuel estimation*, *fuel measurement*

INTRODUCTION

Coastal shrublands dominated by *Baccharis pilularis* (coyote brush) are common in Northern California. This

vegetation type is expanding, due to changes in fire regime, into areas previously dominated by grasses, altering biomass accumulation and potential fire hazard (Russell and McBride 2003). The

precise effect of this conversion is difficult to determine, as there is currently no standard method for estimating biomass in *Baccharis-dominated* communities. The objective of this study was to compare three non-destructive methods of estimating biomass in the coastal *Baccharis* shrub type with the goal of discovering a practical field predictor.

BACKGROUND

Fire is an important factor in the development of *Baccharis* communities. Fire was historically prevalent in this vegetation type, particularly during the Native-American period, and it may serve a role in the regeneration of the dominant species (Schwilk 2002). In addition, the frequency and intensity of fire may be a determining factor in the distribution of *Baccharis* on the landscape. The absence of fire in combination with changes in grazing patterns may be a significant factor in the replacement of grasslands with *Baccharis* shrublands (McBride and Heady, 1968; Elliot and Wehausen, 1974; Williams *et al.*, 1987; Russell and McBride 2003).

Baccharis pilularis is a perennial evergreen shrub common to the coastal areas of California. It is a rapidly growing species especially early in its life cycle. Individual plants may live as long as 100 years, surviving by a process of stem replacement (McBride and Barnhart 2002). Stands in the early stages of succession are characterized by a large number of small shoots. More mature stands have fewer stems which continue to increase in size with age (McBride 1964). On productive sites, *Baccharis* often exceeds two meters in height, creating a significant fuel load.

Comparisons of direct and indirect methods to assess biomass have been

studied in forested vegetation (Reinhardt *et al.*, 2000). Studies have been done regarding biomass in chaparral (Regelbrugge and Conard 1996). However, there are no accepted standardized methods for assessing fuels in coastal shrub vegetation types. In addition, the currently accepted fuel models do not adequately address *Baccharis* and other coastal shrub vegetation (Anderson, 1982; Keane *et al.*, 1994).

Review of previous work on biomass estimation led to the selection of three potential indirect methods for estimating fuel load in the *Baccharis* shrub type. These included height measurement (Russell and McBride 2003), basal stem diameter measurement (Brown 1976), and the estimation of leaf area index (LAI) using a portable hemispheric photography unit, the LAI-2000. Similar tests using LAI have been conducted in forested communities with favorable results (Chen *et al.*, 1991).

We selected as our study areas Golden Gate National Recreation Area and Point Reyes National Seashore (figure 1). These study areas were divided into two distinct ecological types of *Baccharis*, early-successional *Baccharis-dominated* scrub at Point Reyes National Seashore, and mature *Baccharis-dominated* coastal sage scrub at the Marin Headlands at Golden Gate National Recreation Area. The greatest ecological difference between the two sites is that the Point Reyes National Seashore site burned in 1995. The Golden Gate National Recreation Area site had no record of fire. Otherwise, the two parks were similar in terms of vegetation composition, topography, and their proximity to the wildland-urban interface. These sites represent only a small portion of the area being affected by the conversion of grasslands to shrub-

dominated communities, and *Baccharis* represents only one of many shrub types in the region. The need for quantifying biomass in shrub vegetation extends into many different vegetation types. But the methods used to test predictors in this study may be used in any vegetation type.

METHODS

This study was conducted in Point Reyes National Seashore and Golden Gate National Recreation Area in California. Forty-five plots were randomly sampled in each type for a total of 90 plots across the entire study area. Data were collected between May and July in 2002. The data were analyzed for differences both between and within these ecological types. Sample points were randomly located using the Arc View GIS Alaska Tool Pak within polygons designated as *Baccharis* in the Point Reyes National Seashore digital vegetation map (Shirokauer 2001). Only plots with 70-percent *Baccharis* canopy cover were accepted. 70-percent *Baccharis* canopy cover was used to select plots to distinguish the sampling from coastal chaparral community type. If the plot did not meet the criteria, a new direction was chosen at a random azimuth. The plot was then located 10 meters in that direction. If the plot still did not meet the criteria, 90-degrees was added to the first random azimuth. The procedure was repeated at 180-degrees and 270-degrees until an acceptable plot was located. If none of the plots within the 10 meter radius of the original point met the qualifications, the point was not used. All plots were located at least 10 meters away from any trail or road to buffer the measurement from possible edge effects.

LAI using hemispherical photography

The LAI- 2000 (Li-Cor, Inc. Lincoln, NB) was designed for measuring leaf area index for multiple growth forms (grass, shrubs, trees). This instrument has been used primarily to estimate biomass in forest systems (Reinhardt *et al.*, 2000). Measurements made at each of the cardinal directions above and below the canopy were used to determine canopy light interception, from which LAI (leaf area index) was computed using a model of radiative transfer in vegetative canopies (Li-Cor 1992). The standard protocol for shrubs given in the Li-Cor operators' manual was employed, using a 30-degree mask on the sensor to block the operator from reflection on the lens. The LAI estimation was compared with linear regression to biomass results from subsequent clipping and weighing procedures.

Basal stem diameters

Estimation of live fuel biomass from stem diameter measurement using the method described by Brown (1976) was conducted on all sample points. The sample area for this procedure was a one square meter plot centered on the sample point. The diameters of all woody stems were measured five centimeters above the ground to the nearest millimeter. Diameters were converted to basal area and summed for each plot. Basal area estimation was compared using linear regression to biomass results from subsequent clipping and weighing procedures.

Canopy height

Canopy height was estimated at each sample point using a metered pole fixed to the four corners of a 1 square meter PVC plot frame. The base of the plot frame was

placed on the ground with one corner at the sample point and the arms of the plot projecting in the North and East directions. The mean height of the plot was determined by averaging the height at the four stakes. Height was measured to the nearest 0.01-centimeter. Canopy height estimation was compared using linear regression to biomass results from subsequent clipping and weighing procedures.

Biomass

All plant material within each one meter sample plot was removed mechanically. The separated material was then placed in a drying oven at 52-degrees C. Samples were dried until there was no change in weight for three hours. Biomass was measured to the nearest 0.1-gram.

Statistical Analysis

Linear regression models were used to assess the effectiveness of LAI, basal stem diameters, and canopy height in predicting biomass. Normal distribution of residuals was found for variables included in the models. If two or more variables were found to be significantly predictive of biomass, then multiple linear regression models based on these multiple predictors were assessed for any further improvement in predicting biomass. A significance level of 0.05 was used to evaluate the significance of all regression relationships, and adjusted R^2 was used to compare the predictive value between regression models.

RESULTS

Comparison of shrub biomass with three variables indicated a significant linear relationship between crown canopy height and biomass. A weaker but significant linear relationship was also found between basal area and biomass. No linear relationship was found between measured biomass and estimated LAI. Significant differences were found between the two study areas in terms of basal area, height, and biomass, but the relationships between the dependent and independent variables were similar on both study areas.

Linear regression analysis between height and biomass

Linear regression analysis was performed between biomass and height. Several transformations of the biomass variable were tested including, square root, cubic root, and natural log. The residual fit was best with the untransformed variable, and the R^2 changed very little between transformed models, so that none of the transformations were employed in the following analysis. A significant linear relationship was found between biomass (grams) and crown canopy height, (adjusted $R^2 = 0.46$, $P < 0.01$, figure 2). The linear regression equation, biomass (grams) = $238.48 + 3701.36$ height (meters), describes a biomass averaging 238.48 grams (standard error = 43.28 grams) for plots with stakes measuring zero crown canopy height and increasing by 3701.36 grams (standard error = 43.28 grams) per meter increase in height, for each square meter plot.

Linear regression analyses were also performed for each study site individually. Significant relationships were found between biomass and height on both sites.

The linear regression for Golden Gate National Recreation Area was biomass (grams) = $303.20 + 2339.01 \text{ height (meters)}$, with an adjusted $R^2 = 0.33$ and $P < 0.01$. The linear regression equation for Point Reyes National Seashore was biomass (grams) = $950.23 + 4542.73 \text{ height (meters)}$ with an adjusted $R^2 = 0.48$ and $P < 0.01$.

Linear regression analysis between basal area and biomass

A linear regression analysis was performed between biomass and basal area (centimeters²). A significant, but weak, linear relationship was found (adjusted $R^2 = 0.17$, $P < 0.01$). Linear regression analyses were also performed for each study site individually. No significant linear relationship was found between biomass and basal area on Golden Gate National Recreation Area. A significant linear relationship was found between biomass and basal area for Point Reyes National Seashore. The regression equation for Point Reyes National Seashore was biomass (grams) = $51.68 + 0.02 \text{ basal area (centimeters}^2\text{)}$ with an adjusted $R^2 = 0.53$ and $P = 0.01$.

Linear regression analysis between leaf area index (LAI) and biomass

A linear regression analysis was performed for biomass versus estimated LAI. No significant linear relationship was found between estimated LAI and biomass. Linear regression analyses were also performed for each study site individually. No significant linear relationship was found for either site.

Multiple regression analysis between biomass and two independent variables

A multiple regression analysis was conducted with biomass versus height and basal area. A significant relationship was found (adjusted $R^2 = 0.48$, $P < 0.01$). Multiple regression analyses were also performed for each study site individually. Significant linear relationships were found between biomass versus height and basal area on both study sites. The regression for Golden Gate National Recreation Area resulted in an adjusted $R^2 = 0.32$ and $P < 0.01$. The regression statistics for Point Reyes National Seashore were adjusted $R^2 = 0.57$ and $P < 0.01$. A simple linear regression model with biomass (grams) versus a unified variable, height (meters) \times basal area (centimeters²), was also tested. This model indicated a significant linear relationship, but was a relatively weak predictor (adjusted $R^2 = 0.27$, $P < 0.01$).

DISCUSSION

The objective of this study was to compare three nondestructive methods for estimating biomass in coastal *Baccharis* shrublands and to develop a predictive model that can be used in the field for fuel characterization. The three methods tested included crown canopy height measurement, basal area measurement, and leaf area index (LAI) estimation through hemispherical photography. Of these three methods, crown height was the best predictor of measured biomass. Basal area was a weaker predictor, particularly in Golden Gate National Recreation Area, where stands were more mature. The final estimator tested, leaf area index as estimated using the LAI-2000 was not valuable as a predictor.

The linear regression analysis between measured biomass and crown canopy height yielded a useful predictive model. This method was also the most efficient in terms of time and ease of measurement.

Variation occurred in the ratio of height to biomass between the plots sampled on the younger stands in Point Reyes National Seashore compared to the older stands in Golden Gate National Recreation Area, but this variation was much less pronounced than it was for basal stem diameter.

The linear regression analysis between measured biomass and basal area did not yield a useful predictive model. This measure did not predict biomass as well as crown height in part because shrubs that were measured for height and biomass on a specific plot were sometimes rooted outside of that same plot. Using larger plots would have limited this problem, though not eliminated it. The linear regression model for Point Reyes National Seashore was stronger than that for Golden Gate National Recreation Area because the former was a younger stand. Most of the points at Point Reyes were located within the Vision Fire area. This area burned in 1995, eight years prior to our sampling. *Baccharis* responds to fire by adding new shoots so that younger stands of *Baccharis* tend to have a greater number of stems per unit area though the stems are much smaller (McBride 1964). Older stands tend to have fewer, larger stems, and are more likely to be rooted outside of the plot.

Estimated LAI was the least valuable predictor tested. There was no linear relationship found between estimated LAI and measured biomass. The highly layered growth form of *Baccharis* may be one of the reasons for the failure of this technique to predict biomass. The existence of a diverse understory extending into the *Baccharis* canopy may have also contributed to the variability in this method of measurement.

The multiple linear regression model that included both crown canopy height

and basal area versus biomass had the greatest predictive value based on the adjusted R^2 , but it was only slightly better than the model that included crown canopy height alone versus biomass. Since the addition of basal area only marginally increased the predictive power to the regression, the additional effort required to collect basal area data does not seem warranted.

The model developed for this study is applicable only to coastal *Baccharis* communities. However, a similar approach could be tested in other shrub types. As canopy height is the easiest variable to measure, we suggest testing its relationship with biomass in other shrub types in the hope that it will be equally acceptable. It may be that basal area would be a more effective predictor in some shrub types. The data presented in this paper are a first step toward developing predictive tools and fuel models for coastal shrub communities. Further research will be needed to develop complete fuel models for this and other coastal shrub types.

ACKNOWLEDGEMENTS

The authors would like to thank the Joint Fire Science Project and Point Reyes National Seashore for funding this study. The dedicated field crew, including Sayaka Eda, Isaiah Hirschfield, Wende Rehlaender, and Kevin Schwartz, was invaluable. Gary Fellers, Jon Keeley, and Julie Yee provided excellent reviews of this report. We would also like to thank Bob Kean of the Missoula Fire Laboratory for his help with the LAI-2000, and Peggy Herzog, former Fire Ecologist at Point Reyes National Seashore, for helping to develop the project.

REFERENCES

- Anderson, H. 1982. Aids to determining fuel models for estimating fire behavior. USDA Forest Service Intermountain Research Station. Gen. Tech. Rep. INT-122.
- Brown, J.K. 1976. Estimating shrub biomass from basal stem diameters. Canadian Journal of Forest Research 6: 153-158.
- Chen, J. M., T. A. Black, and R. S. Adams. 1991. Evaluation of hemispherical photography for determining Plant Area Index and geometry of forest stands. Agricultural and Forest Meteorology. 56:107-125
- Elliot, H. W. and J. D. Wehausen. 1974. Vegetation succession on coastal rangelands of Point Reyes Peninsula. Madrono 22:231-238.
- Keane, R. E., E. Reinhardt, J.K. Brown. 1994. FOFEM – A First Order Fire Effects Model for predicting the immediate consequences of wildland fire in the United States. In: *Proceedings of the 12th Conference of Fire and Forest Meteorology*. Oct. 26-28, 1993, Jekyll Island, PA. Pp. 628-632.
- Li-Cor. 1992. LAI-2000 Plant Canopy Analyzer: Operating Manual. Li-Cor Inc. Lincoln, Nebraska.
- McBride, J. R., S. Barnhart. 2002. Control of *Baccharis* invasion of grasslands at Mt. Tamalpais State Park. Report to the California State Parks. Unpublished.
- McBride, J. R., H. F. Heady. 1968. Invasion of grassland by *Baccharis pilularis*. Journal of Range Management 21:106-108.
- McBride, J. R. 1964. Ecology of *Baccharis pilularis*. MS Thesis. Department of Forestry. University of California, Berkeley, CA.
- Regelbrugge, J. C., and S. G. Conard. 1996. Biomass and fuel characteristics of chaparral in southern California. *13th Conference on Fire and Forest Meteorology*, Oct 27-31, 1996, Lorne, Australia.
- Reinhardt, E., R. Keane, J. Scott, J.K. Brown. 2000. Quantification of canopy fuels in conifer forests: Assessing crown fuel characteristics using destructive and non-destructive methods. USDA Forest Service Fire Effects Project RWU4403. Unpublished.
- Russell, W. H., J. R. McBride. 2003. Landscape scale vegetation-type conversion and fire hazard in the San Francisco bay area open spaces. Landscape and Urban Planning 64: 201-208.

Shirokauer, D. 2001. Digital vegetation map of Point Reyes National Seashore. U. S. National Park Service, Unpublished.

Schwilk, D. W. 2002. Plant evolution in fire-prone environments. Dissertation: Department of Biological Sciences, Stanford University.

Williams, K., R. J. Hobbs, and S. P. Hamburg. 1987. Invasion of an annual grassland in Northern California by *Baccharis pilularis*. *Oecologia* 72:461-465.



Figure 1. Typical landscape in both sample areas. Point Reyes National Seashore on the left; Golden Gate National Recreation area on the right.




Figure 2. Linear regression, with 95% confidence bands, (adjusted $R^2 = 0.46$), between the crown canopy height of the shrub layer and the dry weight removed from ninety 1 square meter plots within *Baccharis* shrubland in the Point Reyes National Seashore and Golden Gate National Recreation Area, California.