

RESEARCH ARTICLE

MINIMAL PERSISTENCE OF NATIVE BUNCHGRASSES SEVEN YEARS AFTER SEEDING FOLLOWING MASTICATION AND PRESCRIBED FIRE IN SOUTHWESTERN OREGON, USA

Laura M. Busby¹ and Darlene Southworth^{2*}

¹ Bureau of Land Management, Medford District Office,
3040 Biddle Road, Medford, Oregon 97504, USA

² Department of Biology, Southern Oregon University,
1250 Siskiyou Boulevard, Ashland, Oregon 97520, USA

*Corresponding author: Tel.: +1-541-552-6865; e-mail: southworth@sou.edu

ABSTRACT

Seeding of native grasses is widely used to restore plant communities and prevent establishment of introduced species following wildfire and prescribed burns. However, there is a lack of long-term data to evaluate the success of native grass seeding. Here, in the interior valley shrublands of southwestern Oregon, we resurveyed plots that had been masticated and burned, and then seeded with bunchgrasses seven years previously. The prescribed fires had resulted in bare ground that increased opportunities for bunchgrass germination as well as for invasion by introduced plants. After two years, native grass seeding was successful, with increased bunchgrass cover that correlated with decreased cover of introduced species. However, five years later, bunchgrass cover had declined by 80%, and the frequency of plots with bunchgrasses had declined by 60%. Cover of surviving bunchgrasses in year 7 correlated positively with bunchgrass cover in year 2 ($R^2 = 0.34$; $P = 0.003$). Seven years after prescribed fire and seeding, native cover, introduced cover, and species richness were

RESUMEN

La siembra de pastos nativos es ampliamente usada, luego de incendios naturales o de haber realizado quemas prescritas, para restaurar comunidades vegetales y prevenir el restablecimiento de especies introducidas. Sin embargo, la falta de datos a largo plazo impide poder evaluar el éxito de estas siembras. Nosotros re-evaluamos, en el valle de arbustales del interior en el sudoeste de Oregon, EEUU, parcelas que habían sido objeto, siete años antes, de tratamientos de triturado y quemado, y posteriormente sembradas con pastos. Las quemas prescritas habían dejado el suelo desnudo, lo que incrementó las oportunidades de germinación de pastos y también la invasión de especies introducidas. Después de dos años, la siembra de pastos fue exitosa, con un incremento en su cobertura que se correlacionó con un decrecimiento en la cobertura de las especies introducidas. Sin embargo, cinco años después, la cobertura de pastos había declinado en un 80%, y la frecuencia de parcelas con pastos lo habían hecho en un 60%. La cobertura de los pastos sobrevivientes se correlacionó positivamente con la cobertura en el año 2 ($R^2 = 0.34$; $P = 0.003$). Después de siete años de las quemas prescritas y el sembrado, la cobertura de especies nativas, introducidas, y la riqueza de especies

unchanged, and bunchgrass persistence was minimal. Basically, seeding following mastication and prescribed burning had a minimal effect. This study highlights the importance of longer-term monitoring to determine the efficacy of seeding treatments.

permaneció invariable, y la persistencia de pastos fue mínima. Básicamente, la siembra de pastos luego de los tratamientos de triturado y de quemas tuvo un efecto mínimo. Este estudio subraya la importancia del monitoreo a largo plazo para determinar la eficacia de los tratamientos de siembra.

Keywords: *Achnatherum lemmonii*, *Bromus carinatus*, *Elymus glaucus*, *Festuca roemerii* var. *klamathensis*, long-term monitoring, restoration ecology, southern Oregon chaparral

Citation: Busby, L., and D. Southworth. 2014. Minimal persistence of native bunchgrasses seven years after seeding, following mastication and prescribed fire in southwestern Oregon, USA. *Fire Ecology* 10(3): 63–71. doi: 10.4996/fireecology.1003063

INTRODUCTION

Seeding of native grasses is widely used to restore particular species, plant communities, and ecosystem services following wildfire and prescribed burns (e.g., Hardegree *et al.* 2011). However, there is a lack of long-term data to evaluate the success of native grass establishment and its effect in preventing introduced species invasions (Robichaud *et al.* 2009, Peppin *et al.* 2010, Endress *et al.* 2012). Such studies are needed to evaluate the benefits of restoration projects and to improve restoration techniques (Duren and Muir 2010, Hardegree *et al.* 2011, Peppin *et al.* 2011).

Invasion by introduced species is a threat to the conservation of shrublands and creates a challenge to restoration (Hosten *et al.* 2006, Endress *et al.* 2012). Both mastication and fire lead to more introduced species in part because of increased bare ground (Beyers 2004, Potts and Stephens 2009). At a site with a Mediterranean climate (Hierro *et al.* 2006), prescribed fire created a favorable environment for invasion by yellow star-thistle (*Centaurea solstitialis* L.). However, in a ponderosa pine (*Pinus ponderosa* Lawson and C. Lawson) forest, the combination of mastication plus burning resulted in greater native plant diversity with only a slight increase in introduced species (Kane *et al.* 2010).

Seeding with native plants may be a way to prevent the spread of introduced species, but a review of post-fire seeding studies found equivalent numbers of effective and ineffective results on introduced species (Peppin *et al.* 2010). In a grassland-savannah ecosystem, post-fire seeding with native species reduced establishment of introduced species and increased native plant cover and species richness; however, the benefits of seeding occurred only at intermediate levels of introduced species richness (Suding and Gross 2006).

In dense, fire-suppressed shrublands that had been masticated, Coulter *et al.* (2010) seeded native bunchgrass species following prescribed burns. Post-fire seeding and subsequent increased bunchgrass cover correlated with decreased introduced species. Bunchgrasses were not present on unseeded plots. Initially, the native grass seeding appeared to be successful with 10% cover of seeded species after two growing seasons. After five more years, we resurveyed their plots to determine the longer-term success of native grass seeding treatments.

Our objective was to assess the longer-term effectiveness of seeding with native bunchgrasses following brush mastication and prescribed fire in the shrublands of an interior valley in southern Oregon, USA. We compared the cover of bunchgrasses and intro-

duced species on masticated plots resurveyed seven years after the prescribed burn with the cover two years after the prescribed burn. We hypothesized that (1) bunchgrasses that had survived two growing seasons would persist and expand; (2) greater bunchgrass cover in year 2 would predict cover in year 7; and (3) introduced species cover would be lower on plots with greater bunchgrass cover. In addition, we asked whether the treatments of prescribed fire and bunchgrass seeding would change the plant community.

METHODS

We resurveyed the sites used by Coulter *et al.* (2010). Sites were on south facing slopes in the Applegate Valley near Ruch, Oregon, USA: China Gulch (42° 15' N, 122° 3' W) and Hukill Hollow (42° 11' N, 122° 59' W). The climate is a Mediterranean type with cool, wet winters and hot, dry summers: mean temperature 4°C in January and 21°C in July. For the years 2005 to 2012, annual rainfall ranged from 343 mm to 734 mm, with only 10% of total precipitation in the summer months, June through September (www.wrcc.dri.edu/cgi-bin/cliMONtpr.pl?or7391, accessed 20 Oct 2014). Summer rainfall varied from 0.5 cm to 7 cm during the interval between surveys (Figure 1).




Figure 1. Summer rainfall (June through September) at the Ruch, Oregon, USA, gauging station. Between the second growing season (2007) and the time of resurvey (2012), there was an exceedingly dry summer (2008).

Prior to mastication, the sites were dominated by buckbrush (*Ceanothus cuneatus* [Hook.] Nutt.), sticky whiteleaf manzanita (*Arctostaphylos viscida* Parry), Pacific poison oak (*Toxicodendron diversilobum* [Torr. & A. Gray] Greene), and California yerba santa (*Eriodictyon californicum* [Hook. & Arn.] Torr.). Mastication was employed to reduce the threat of high-severity wildfire by lowering fuels from standing vegetation to ground level (Kane *et al.* 2009, Southworth *et al.* 2011, Duran *et al.* 2012). Five years after mastication, the sites were treated with prescribed fire to reduce woody fuels on the ground.

At each of two sites, China Gulch (CG) masticated in 2001 and Hukill Hollow (HH) masticated in 2002, 30 1 m² plots were installed randomly and the vegetation surveyed (Coulter *et al.* 2010). All plots (60 total) were treated with a prescribed burn in October 2005, year 0. Two weeks after the burn, 15 plots at each site (30 total) were seeded with four native bunchgrasses: Lemmon's needlegrass (*Achnatherum lemmonii* [Vasey] Barkworth, syn. *Stipa lemmonii* [Vasey] Scribn.), California brome (*Bromus carinatus* Hook. & Arn.), blue wildrye (*Elymus glaucus* Buckley), and Roemer's fescue (*Festuca roemeri* [Pavlick] S. Aiken var. *klamathensis* B.L. Wilson, syn. *F. idahoensis* Elmer sp. *roemeri* [Pavlick] S. Aiken). An equal number of paired control plots (30 total) were unseeded. Cover of all species was assessed in the first growing season, year 1 (2006), and in year 2 (2007). In June 2012, year 7, we resurveyed the seeded ($n = 30$) and unseeded ($n = 30$) plots. We estimated percent cover of plant species using midpoint values of FIREMON classes (0% to 1%, >1% to 5%, >5% to 15%, >15% to 25%, >25% to 35%, >35% to 45%, >45% to 55%, >55% to 65%, >65% to 75%, >75% to 85%, >85% to 95%, and >95% to 100%) (http://frames.nbio.gov/projects/firemon/FIREMON_SamplingMethods.pdf).

We used a two sample *t*-test in MINITAB v. 15 (Minitab Inc., State College, Pennsylvania).

nia, USA) with a significance threshold of $\alpha = 0.05$ to compare bunchgrass cover in seeded and unseeded plots at replicate sites and to evaluate the change in species richness and percent cover from year 2 to year 7. We analyzed changes in percent cover of species within the same plot using paired t -tests, and changes in the frequency of plots with bunchgrasses in different years using χ^2 . We performed a linear regression to compare the relationship between bunchgrass and introduced species cover.

To determine differences between the plant communities of seeded and unseeded plots, we compared seeded and unseeded groups by multi-response permutation procedures (MRPP) in PC-ORD v. 6 using Sørensen (Bray-Curtis) distance measures (McCune and Grace 2002, Peck 2010). We calculated A , the chance-corrected within-group agreement, and the P -value. We analyzed the two sites separately because they differed in total number of species.

RESULTS

Total plant alpha-diversity was 58 species at two sites (Appendix 1). Of these, 31% were introduced species; 22% were grasses. Beyond the seeded grasses, only one other grass species was native. Four shrub species—buckbrush, California yerba santa, pink

honeysuckle (*Lonicera hispidula* [Lindl.] Douglas ex Torr. & A. Gray), and Pacific poison oak—were present in year 7.

On unseeded plots, bunchgrass cover was 0.4%. On seeded plots, bunchgrass cover was 1.9% in year 7, a decline in cover from 9.7% in year 2 ($t = 3.01$; $P = 0.005$; Table 1). California brome and blue wildrye declined from year 2 to year 7 with cover losses of 91% to 96%, while cover of Lemmon's needlegrass and Roemer's fescue remained the same. All four seeded bunchgrass species remained at low cover levels (0.1% to 0.9%) and each was found on at least one plot.

Since bunchgrasses had not germinated in many plots, we looked closely at the set of 23 plots (21 seeded, 2 unseeded) in which bunchgrasses had been present by year 2 (Table 2). The total number of plots with bunchgrasses declined from 23 to 9 (7 seeded, 2 unseeded) in year 7 ($\chi^2 = 7.2$, $P = 0.007$). Between year 2 and year 7, cover declined in California brome ($t = 4.10$, $P = 0.001$) and in blue wildrye ($t = 3.08$, $P = 0.006$) but did not change in Roemer's fescue ($t = -0.08$, $P = 0.94$) or in Lemmon's needlegrass ($t = 3.08$, $P = 0.50$) (Table 2). Bunchgrasses with higher cover by year 2 persisted into year 7 better than those with less cover ($R^2 = 0.34$; $P = 0.003$; Figure 2).

Introduced plant cover increased in both seeded and unseeded plots over the seven-year period (Table 3). However, changes in intro-

Table 1. Bunchgrass percent cover (mean) on seeded ($n = 30$) and unseeded ($n = 30$) 1 m² plots before and after fall burns of masticated shrublands at China Gulch and Hukill Hollow, in southern Oregon, USA. Compare with Table 3 in Coulter *et al.* (2010).

Bunchgrass	Percent cover						
	Year 0	Year 1		Year 2		Year 7	
		Unseeded	Seeded	Unseeded	Seeded	Unseeded	Seeded
Lemmon's needlegrass	0.0	0.0	0.1	0.0	0.3	0.0	0.2
California brome	0.8	0.2	3.5	0.2	2.7	0.0	0.1
Blue wildrye	0.0	0.0	0.7	0.0	5.9	0.4	0.7
Roemer's fescue	0.0	0.0	0.9	0.0	0.8	0.0	0.9
Total bunchgrasses	0.8	0.2	5.2	0.2	9.7	0.4	1.9

Table 2. Changes in number of plots ($n = 60$; 30 seeded, 30 unseeded) with surviving bunchgrasses, between year 2 and year 7 at China Gulch and Hukill Hollow, in southern Oregon, USA. Bunchgrass cover (%) is the mean cover for each surviving species on only the plots with that species.

Bunchgrass	Plots with bunchgrass year 2		Plots with bunchgrass year 7	
	Number	Cover (%)	Number	Cover (%)
Lemmon's needlegrass	5	2.0	2	3.0
California brome	16	5.5	1	3.0
Blue wildrye	19	9.3	6	5.7
Roemer's fescue	6	4.2	3	8.7




Figure 2. Correlation between bunchgrass cover in year 7 with that in year 2 on plots with bunchgrass survival, at China Gulch and Hukill Hollow in southern Oregon, USA.

duced cover between seeded and unseeded plots did not differ ($P = 0.68$). Introduced species richness increased in both seeded and unseeded plots. Species richness increased by an average of two species. There was no difference between seeded and unseeded plots in the

percent increase of introduced species richness ($P = 0.73$). In plots with bunchgrasses present in year 7, introduced cover had not changed from pre-burn cover levels although there had been a substantial increase in year 2 (Figure 3; $R^2 = 0.05$, $P = 0.61$).

Native plant cover decreased in both seeded and unseeded plots over the seven-year period (Table 3). Changes in native cover did not differ between seeded and unseeded plots ($P = 0.46$). Native species richness increased in both seeded and unseeded plots with species richness increasing by an average of one species. Seeded and unseeded plots did not differ in percent increase of native species richness ($P = 0.81$).

In year 7, plant communities on seeded and unseeded plots did not differ at either CG ($A = -0.004$, $P = 0.62$) or at HH ($A = 0.0126$, $P = 0.08$). This was consistent with the lack of differences in introduced or native species between seeded and unseeded plots.

Table 3. Change in cover and species richness of native and introduced plants as the differences in total cover (Δ cover) and in the fraction of total cover (% change) from year 0 to year 7 at China Gulch and Hukill Hollow, in southern Oregon, USA. Changes in percent cover (mean) and species richness (mean) were based on 1 m² plots, seeded and unseeded ($n = 30$ each). Differences between unseeded and seeded plots were not significant. Compare with Table 2 in Coulter *et al.* (2010).

Treatment	Cover				Species richness			
	Native		Introduced		Native		Introduced	
	Δ cover	% change	Δ cover	% change	Δ species	% change	Δ species	% change
Unseeded	-13.8	-19.4	10.8	41.4	0.9	15.5	2.0	85.9
Seeded	-19.4	-27.2	14.2	62.7	1.0	18.0	1.9	83.6




Figure 3. Cover of introduced species in years 0, 1, 2, and 7 on all seeded (solid line) and unseeded (dashed line) plots at China Gulch and Hukill Hollow in southern Oregon, USA. Seeded and unseeded treatments did not differ.

DISCUSSION

Hypothesis 1, that bunchgrasses that had survived two growing seasons would persist and expand, was not supported. Seeded bunchgrasses that were present two years following prescribed fire and seeding persisted poorly after seven years, with low bunchgrass cover occurring on only a few plots. Although bunchgrass cover in year 2 was nearly 10% following fall burns of brush-masticated shrublands, bunchgrass cover declined to less than 2% in year 7. While all four species of bunchgrasses remained on at least one plot, all cover estimates were low. Relative survival of the four seeded bunchgrass species differed. Lemmon's needlegrass and Roemer's fescue persisted with low cover while California brome and blue wildrye attained higher cover in year 2 and then declined by year 7.

Several factors may have prevented successful establishment of seeded bunchgrasses. One, our sites were on southerly aspects with a high concomitant heat load. In California, planted grasses established better on north facing than on south facing slopes (Lulow *et al.* 2007). Two, rainfall at our sites varied annually over the seven-year course of the study and included one exceedingly dry year (2008),

which may have created conditions in which only well-established plants survived. Three, because California brome and blue wildrye are short-lived perennials in contrast to longer lived Roemer's fescue and Lemmon's needlegrass (<http://plants.usda.gov>), they may have reached the ends of their lifespans and failed to reseed due to site conditions or the drought year.

Supporting hypothesis 2, bunchgrass cover in year 7 correlated positively with cover in year 2. Greater cover may indicate better root development, better competitive ability, and consequently longer persistence. In other studies, larger bunchgrass transplants showed better survival and had greater final mass (Wilson 1994, Page and Bork 2005).

Hypothesis 3, that introduced species cover would be lower on plots with greater bunchgrass cover, was not supported. Seeding with bunchgrasses had no effect on the cover or species richness of introduced or native plant species. Although the cover of native and introduced species had decreased slightly and species richness had increased slightly after two years, after seven years neither cover nor species richness of native and introduced species differed between seeded and unseeded plots. Introduced species cover showed no correlation with bunchgrass cover. Similarly, neither richness nor cover of native species was affected by seeding with bunchgrasses. While native grass seeding limited the invasion of introduced grasses in a grassland-savanna ecosystem (Endress *et al.* 2012), a review of post-fire seeding studies found equivalent numbers of effective and ineffective results of seeding on introduced species (Peppin *et al.* 2010).

Treatments of prescribed fire and bunchgrass seeding did not change the plant community, due to the stability of both native and introduced components of vegetation. Apparently, species adapted to seasonally dry habitats through characteristics such as below-ground perennating structures and seed banks persist through prescribed fire.

These findings show the importance of longer-term monitoring of restoration treatments to determine their success (Robichaud *et al.* 2009, Peppin *et al.* 2010, Endress *et al.* 2012). Initial results at our study sites indicated a successful introduction of native bunchgrasses, but this introduction was considerably less successful after seven years. The outcome of seeding may depend on habitat conditions and on climatic variability. Although the trend has been to seed with native species on public lands to meet natural ecological system objectives, the success of such treatments remains uncertain (Peppin *et al.* 2011).

Climate change is likely to interfere with attempts at restoration of native plant communities, particularly at lower elevations and on south facing slopes with little summer rainfall (Harris *et al.* 2006, Batchelet *et al.* 2011). The shrubland communities of inland valleys may be adapted to Mediterranean-type climates, and thus will survive and expand. However, our ability to manage them or to promote native species restoration may be more limited. If reseeding is not effective, preservation of habitat becomes even more important.

ACKNOWLEDGEMENTS

We thank C.T. Coulter for help in relocating plots and in standardizing data collection protocols, D. Kendig for project support, and S. Joachims for field assistance. Research was supported in part by the US Department of Interior, Bureau of Land Management, Medford District. The views and conclusions contained in this document are those of the authors and should not be interpreted as representing the opinions or policies of the US government. Mention of trade names or commercial products does not constitute their endorsement by the US government.

LITERATURE CITED

- Bachelet, D., B.R. Johnson, S.D. Bridgham, P.V. Dunn, H.E. Anderson, and B.M. Rogers. 2011. Climate change impacts on western Pacific Northwest prairies and savannas. *Northwest Science* 85: 411–429. doi: [10.3955/046.085.0224](https://doi.org/10.3955/046.085.0224)
- Beyers, J.L. 2004. Postfire seeding for erosion control: effectiveness and impacts on native plant communities. *Conservation Biology* 18: 947–956. doi: [10.1111/j.1523-1739.2004.00523.x](https://doi.org/10.1111/j.1523-1739.2004.00523.x)
- Coulter, C.T., P.E. Hosten, and D. Southworth. 2010. Prescribed fire and post-fire seeding in brush masticated oak-chaparral: consequences for native and non-native plants. *Fire Ecology* 6(2): 60–75. doi: [10.4996/fireecology.0602060](https://doi.org/10.4996/fireecology.0602060)
- Duren, O.C., and P.S. Muir. 2010. Does fuels management accomplish restoration in southwest Oregon, USA, chaparral? Insights from age structure. *Fire Ecology* 6(2): 76–96. doi: [10.4996/fireecology.0602076](https://doi.org/10.4996/fireecology.0602076)
- Duren, O.C., P.S. Muir, and P.E. Hosten. 2012. Vegetation change from the Euro-American settlement era to the present in relation to environment and disturbance in southwest Oregon. *Northwest Science* 86: 310–328. doi: [10.3955/046.086.0407](https://doi.org/10.3955/046.086.0407)
- Endress, B.A., C.G. Parks, B.J. Naylor, S.R. Radosevich, and M. Porter. 2012. Grassland response to herbicides and seeding of native grasses 6 years posttreatment. *Invasive Plant Science and Management* 5: 311–316. doi: [10.1614/IPSM-D-11-00050.1](https://doi.org/10.1614/IPSM-D-11-00050.1)
- Hardegree, S.P., T.A. Jones, B.A. Roundy, M.L. Shaw, and T.A. Monaco. 2011. Assessment of range planting as a conservation practice. Pages 171–212 in: D.D. Briske, editor. *Conservation benefits of rangeland practices: assessment, recommendations, and knowledge gaps*. Allen Press, Lawrence, Kansas, USA.

- Harris, J.A., R.J. Hobbs, E. Higgs, and J. Aronson. 2006. Ecological restoration and global climate change. *Restoration Ecology* 14: 170–176. doi: [10.1111/j.1526-100X.2006.00136.x](https://doi.org/10.1111/j.1526-100X.2006.00136.x)
- Hierro, J.L., D. Villarreal, O. Eren, J.M. Graham, and R.M. Callaway. 2006. Disturbance facilitates invasion: the effects are stronger abroad than at home. *American Naturalist* 168: 144–156. doi: [10.1086/505767](https://doi.org/10.1086/505767)
- Hosten, P.E., G. Hickman, F. Lake, F. Lang, and D. Vesely. 2006. Oak woodland and savanna restoration. Pages 63–96 in: D. Apostol and M. Sinclair, editors. *Restoring the Pacific Northwest: the art and science of ecological restoration in Cascadia*. Island Press, Washington, D.C., USA.
- Kane, J.M., J.M. Varner, and E.E. Knapp. 2009. Novel fuelbed characteristics associated with mechanical mastication treatments in northern California and southwestern Oregon. *International Journal of Wildland Fire* 18: 686–697. doi: [10.1071/WF08072](https://doi.org/10.1071/WF08072)
- Kane, J.M., J.M. Varner, E.E. Knapp, and R.F. Powers. 2010. Understory vegetation response to mechanical mastication and other fuels treatments in a ponderosa pine forest. *Applied Vegetation Science* 13: 207–220. doi: [10.1111/j.1654-109X.2009.01062.x](https://doi.org/10.1111/j.1654-109X.2009.01062.x)
- Lulow, M.E., T.P. Young, J.L. Wirka, and J.H. Anderson. 2007. Variation in the initial success of seeded native bunchgrasses in the rangeland foothills of Yolo County, California. *Ecological Restoration* 25: 20–28. doi: [10.3368/er.25.1.20](https://doi.org/10.3368/er.25.1.20)
- McCune, B., and J.B. Grace. 2002. *Analysis of ecological communities*. MjM Software Design, Gleneden Beach, Oregon, USA.
- Page, H.N., and E.W. Bork. 2005. Effect of planting season, bunchgrass species, and neighbor control on the success of transplants for grassland restoration. *Restoration Ecology* 13: 651–685. doi: [10.1111/j.1526-100X.2005.00083.x](https://doi.org/10.1111/j.1526-100X.2005.00083.x)
- Peck, J.E. 2010. *Multivariate analysis for community ecologists: step-by-step using PC-ORD*. MjM Software Design, Gleneden Beach, Oregon, USA.
- Peppin, D.L., P.Z. Fulé, C.H. Sieg, J.L. Beyers, and M.E. Hunter. 2010. Post-wildfire seeding in forest of the western United States: an evidence-based review. *Forest Ecology and Management* 260: 573–586. doi: [10.1016/j.foreco.2010.06.004](https://doi.org/10.1016/j.foreco.2010.06.004)
- Peppin, D.L., P.Z. Fulé, C.H. Sieg, J.L. Beyers, M.E. Hunter, and P. Robichaud. 2011. Recent trends in post-wildfire seeding in western US forests: costs and seed mixes. *International Journal of Wildland Fire* 20: 702–708. doi: [10.1071/WF10044](https://doi.org/10.1071/WF10044)
- Potts, J.B., and S.L. Stephens. 2009. Invasive and native plant responses to shrubland fuel reduction: comparing prescribed fire, mastication, and treatment season. *Biological Conservation* 142: 1657–1664. doi: [10.1016/j.biocon.2009.03.001](https://doi.org/10.1016/j.biocon.2009.03.001)
- Robichaud, P.R., S.A. Lewis, R.E. Brown, and L.E. Ashmun. 2009. Emergency post-fire rehabilitation treatment effects on burned area ecology and long-term restoration. *Fire Ecology* 5(1): 115–128. doi: [10.4996/fireecology.0501115](https://doi.org/10.4996/fireecology.0501115)
- Southworth, D., J. Donohue, J.L. Frank, and J. Gibson. 2011. Mechanical mastication and prescribed fire in conifer-hardwood chaparral: differing responses of ectomycorrhizae and truffles. *International Journal of Wildland Fire* 20: 888–896. doi: [10.1071/WF10033](https://doi.org/10.1071/WF10033)
- Suding, K.N., and K.L. Gross. 2006. Modifying native and non-native species richness correlations: the influence of fire and seed addition. *Ecological Applications* 16: 1319–1326. doi: [10.1890/1051-0761\(2006\)016\[1319:MNAESR\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[1319:MNAESR]2.0.CO;2)
- Wilson, S.D. 1994. Initial size and the competitive responses of two grasses at two levels of soil nitrogen: a field experiment. *Canadian Journal of Botany* 72: 1349–1354. doi: [10.1139/b94-165](https://doi.org/10.1139/b94-165)

Appendix 1. Species present at China Gulch (CG) and Hukill Hollow (HH) in the Applegate Valley, Oregon, USA, in year 7. Species were characterized by life form (seeded bunchgrass = bg; forb = f; graminoid = g; woody shrub = s) and origin (introduced = I; native = N). Cover was assessed as mean percent cover on 30 1 m² plots at each site.

Taxon	Common name	Life form	Origin	% cover	
				CG	HH
<i>Achillea millefolium</i> L.	common yarrow	f	N	0.0	0.3
<i>Achnatherum lemmonii</i> (Vasey) Barkworth	Lemmon's needlegrass	bg	N	0.0	0.2
<i>Agoseris</i> sp.	agoseris	f	N	0.0	0.7
<i>Aira caryophylla</i> L.	silver hairgrass	g	I	0.0	3.1
<i>Amsinckia menziesii</i> (Lehm.) A. Nelson & J.F. Macbr.	Menzies' fiddleneck	f	N	0.0	0.3
<i>Aster</i> spp.	aster	f	N	0.1	1.7
<i>Avena fatua</i> L.	wild oat	g	I	11.2	0.0
<i>Bromus carinatus</i> Hook. & Arn.	California brome	bg	N	0.0	0.1
<i>Bromus diandrus</i> Roth	rippgut brome	g	I	0.1	0.0
<i>Bromus hordeaceus</i> L.	soft brome	g	I	8.6	4.3
<i>Bromus madritensis</i> L.	compact brome	g	I	3.3	0.0
<i>Bromus tectorum</i> L.	cheat grass	g	I	1.9	1.4
<i>Calystegia occidentalis</i> (A. Gray) Brummitt	chaparral false bindweed	f	N	0.1	0.0
<i>Castilleja tenuis</i> (A. Heller) T.I. Chuang & Heckard	hairy Indian paintbrush	f	N	0.0	0.2
<i>Ceanothus cuneatus</i> (Hook.) Nutt.	buckbrush	s	N	0.0	1.3
<i>Centaurea solstitialis</i> L.	yellow star-thistle	f	I	0.0	0.7
<i>Cerastium arvense</i> L.	field chickweed	f	N	0.0	0.4
<i>Cirsium cymosum</i> (Greene) J.T. Howell	peregrine thistle	f	N	5.8	0.0
<i>Clarkia purpurea</i> (W. Curtis) A. Nelson & J.F. Macbr. ssp. <i>quadrivulnera</i> (Douglas ex Lindl.) F.H. Lewis & M.I. Lewis	winecup clarkia	f	N	2.3	3.3
<i>Clarkia rhomboidea</i> Douglas ex Hook.	diamond clarkia	f	N	0.0	0.2
<i>Collomia grandiflora</i> Douglas ex Lindl.	grand collomia	f	N	0.0	0.2
<i>Cryptantha flaccida</i> (Douglas ex Lehm.) Greene	weakstem cryptantha	f	N	1.7	0.0
<i>Cynosurus echinatus</i> L.	bristly dogstail grass	f	I	0.0	0.9
<i>Daucus pusillus</i> Michx.	American wild carrot	f	N	2.3	4.8
<i>Dichelostemma congestum</i> (Sm.) Kunth	ookow	f	N	0.1	0.0
<i>Elymus glaucus</i> Buckley	blue wildrye	bg	N	0.4	0.8
<i>Eriodictyon californicum</i> (Hook. & Arn.) Torr.	California yerba santa	s	N	5.6	0.1
<i>Erodium cicutarium</i> (L.) L'Hér. ex Aiton	redstem stork's bill	f	I	0.0	1.5
<i>Eriophyllum lanatum</i> (Pursh) Forbes	common woolly sunflower	f	N	0.2	1.2
<i>Festuca roemerii</i> (Pavlick) S. Aiken var. <i>klamathensis</i> B.L. Wilson	Roemer's fescue	bg	N	0.1	0.8
<i>Galium ambiguum</i> W. Wight	Yolla Bolly bedstraw	f	N	0.0	0.1
<i>Galium aparine</i> L.	stickywilly	f	N	0.0	0.0
<i>Galium parisiense</i> L.	wall bedstraw	f	I	0.0	1.6
<i>Galium</i> sp.	bedstraw	f	N	0.0	0.0
<i>Gilia capitata</i> Sims	bluehead gilia	f	N	0.2	0.0
<i>Hieracium</i> sp.	hawkweed	f	N	0.0	4.2
<i>Hypericum perforatum</i> L.	common St. Johnswort	f	I	0.0	0.0
<i>Lactuca serriola</i> L.	lettuce	f	I	0.0	0.4
<i>Leptosiphon bicolor</i> Nutt.	true babystars	f	N	0.0	1.9
<i>Lomatium californicum</i> (Nutt.) Mathias & Constance	California lomatium	f	N	0.0	0.1
<i>Lomatium macrocarpum</i> (Nutt. ex Torr. & A. Gray) J.M. Coult. & Rose	bigseed biscuitroot	f	N	0.0	0.1
<i>Lonicera hispidula</i> (Lindl.) Douglas ex Torr. & A. Gray	pink honeysuckle	s	N	0.0	0.2
<i>Lotus micranthus</i> Benth.	desert deervetch	f	N	0.0	1.9
<i>Madia citriodora</i> Greene	lemonscented madia	f	N	0.0	0.7
<i>Madia gracilis</i> (Sm.) D.D. Keck	grassy tarweed	f	N	14.8	10.4
<i>Madia madioides</i> (Nutt.) Greene	woodland madia	f	N	0.1	0.9
<i>Micropus californicus</i> Fisch. & C.A. Mey.	q-tips	f	N	0.0	1.8
<i>Phacelia heterophylla</i> Pursh	varileaf phacelia	f	N	0.0	0.5
<i>Plagiobothrys nothofulvus</i> (A. Gray) A. Gray	rusty popcornflower	f	N	0.2	0.0
<i>Plectritis congesta</i> (Lindl.) DC.	shortspur seablush	f	N	0.0	0.2
<i>Poa bulbosa</i> L.	bulbous bluegrass	g	I	0.0	0.0
<i>Sisymbrium altissimum</i> L.	tall tumbledustard	f	I	0.0	0.1
<i>Stellaria media</i> (L.) Vill.	common chickweed	f	I	0.0	0.3
<i>Torilis arvensis</i> (Huds.) Link	spreading hedgeparsley	f	I	4.9	2.7
<i>Toxicodendron diversilobum</i> (Torr. & A. Gray) Greene	Pacific poison oak	s	N	6.2	15.8
<i>Tragopogon dubius</i> Scop.	yellow salsify	f	I	0.0	0.0
<i>Vulpia microstachys</i> (Nutt.) Munro	small fescue	g	N	0.8	0.1
<i>Vulpia myuros</i> (L.) C.C. Gmel.	annual fescue	g	I	6.6	0.0